

AD-A193 615

STRONG LAW FOR MIXING SEQUENCE(U) PITTSBURGH UNIV PA  
CENTER FOR MULTIVARIATE ANALYSIS X CHEN ET AL. DEC 87  
TR-84-47 AFOSR-TR-88-0401 F49620-85-C-0008


1/1

UNCLASSIFIED

F/G 12/3

NL





MICROCOPY RESOLUTION TEST CHART  
 (JREAU) U.S. STANDARDS-1963-A

AD-A193 615

DTIC FILE COPY

②

AFOSR-TR- 88-0401


STRONG LAW FOR MIXING SEQUENCE\*

Xiru Chen and Yuehua Wu

Center for Multivariate Analysis  
University of Pittsburgh

Technical Report No. 87-47

**Center for Multivariate Analysis**  
**University of Pittsburgh**



DTIC  
ELECTE  
MAY 03 1988  
S H D

DISTRIBUTION STATEMENT A

Approved for public release  
Distribution Unlimited

88 5 02 284

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER <b>AFOSR-TR-88-0401</b>	2. GOVT ACCESSION NO. <b>A193615</b>	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) <b>Strong law for mixing sequence</b>		5. TYPE OF REPORT & PERIOD COVERED <b>Journal Technical - December 1987</b>
7. AUTHOR(s) <b>Xiru Chen and Yuehua Wu</b>		6. PERFORMING ORG. REPORT NUMBER <b>87-47</b>
9. PERFORMING ORGANIZATION NAME AND ADDRESS <b>Center for Multivariate Analysis Fifth Floor Thackeray Hall University of Pittsburgh, Pittsburgh, PA 15260</b>		8. CONTRACT OR GRANT NUMBER(s) <b>F49620-85-C-0008</b>
11. CONTROLLING OFFICE NAME AND ADDRESS <b>Air Force Office of Scientific Research Department of the Air Force Bolling Air Force Base, DC 20332</b>		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS <b>61102F 330446</b>
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) <b>AFOSR BK1410 BAF B DC 20332-6448</b> <i>nm</i>		12. REPORT DATE <b>December 1987</b>
		13. NUMBER OF PAGES <b>13</b>
		15. SECURITY CLASS. (of this report) <b>Unclassified</b>
16. DISTRIBUTION STATEMENT (of this Report)  <b>Approved for public release; distribution unlimited.</b>		18a. DECLASSIFICATION/DOWNGRADING SCHEDULE
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) <b>mixing coefficient; stationary sequence; strong law of large numbers.</b>		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) <b>In this note we present some theorems on the strong law for the mixing sequence which is not necessarily stationary, and the mixing coefficient involving only a pair of variables in the sequence.</b>		

DD FORM 1 JAN 73 1473

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

1000-00-27-82000

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

2

STRONG LAW FOR MIXING SEQUENCE\*

Xiru Chen and Yuehua Wu

Center for Multivariate Analysis  
University of Pittsburgh

Technical Report No. 87-47

December 1987

Center for Multivariate Analysis  
Fifth Floor Thackeray Hall  
University of Pittsburgh  
Pittsburgh, PA 15260

DTIC  
ELECTE  
S MAY 03 1988 D  
H

\* Research sponsored by the Air Force Office of Scientific Research under Contract F49620-85-C-0008. The United States Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon.

**DISTRIBUTION STATEMENT A**

Approved for public release;  
Distribution Unlimited

# STRONG LAW FOR MIXING SEQUENCE \*

Xiru Chen and Yuehua Wu

## ABSTRACT

In this note we present some theorems on the strong law for the mixing sequence which is not necessarily stationary, and the mixing coefficient involving only a pair of variables in the sequence.

*AMS 1980 Subject Classifications:* Primary 60F15.

*Key words and phrases:* mixing coefficient, stationary sequence, strong law of large numbers.

\* Research sponsored by the Air Force Office of Scientific Research under Contract F49620-85-C-0008. The United States Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon.

## 1. INTRODUCTION

In this article we present some results concerning the strong law of a mixing sequence  $\{X_n, n \geq 1\}$ . We do not assume that  $\{X_n\}$  is stationary, and we use mixing coefficients involving only a pair of variables  $X, Y$  (in that order): The Rosenblatt mixing coefficient

$$\alpha(X, Y) = \sup\{|P(X \in A, Y \in B) - P(X \in A)P(Y \in B)| : A \in B', B \in B'\}$$

and the Ibragimov mixing coefficient

$$\beta(X, Y) = \sup\{|P(Y \in B | X \in A) - P(Y \in B)| : A \in B', B \in B', P(X \in A) > 0\}$$

where  $B'$  is the  $\sigma$ -field of all Borel sets in  $R'$ .

THEOREM 1. Suppose that  $\{X_n, n \geq 1\}$  is a sequence of random variables, and for some  $p > 1$  the following conditions are satisfied:

$$1^\circ. \sup_n E|X_n|^p < \infty. \quad (1)$$

$$2^\circ. \text{ There exists } \epsilon > 0 \text{ such that as } |i - j| \rightarrow \infty,$$

$$\alpha(X_i, X_j) \leq \rho(|i - j|) = \begin{cases} O(|i - j|^{-p/(2p-2)-\epsilon}), & 1 < p < 2, \\ O(|i - j|^{-2/p-\epsilon}), & p \geq 2. \end{cases} \quad (2)$$

Then

$$\lim_{n \rightarrow \infty} (S_n - ES_n)/n = 0, \quad \text{a.s.} \quad (3)$$

Here and in the sequel  $S_n = \sum_{i=1}^n X_i$ .

THEOREM 2. Suppose that  $\{X_n, n \geq 1\}$  is a sequence of random variables, and one of the following conditions are satisfied:

$$(I) \quad \sum_{n=1}^{\infty} \text{var}(X_n)/n^2 < \infty, \quad \sup_n E|X_n| < \infty,$$

and

$$\beta(X_i, X_j) \leq \mu(|i - j|), \quad \sum_{n=0}^{\infty} \mu^{1/2}(n) < \infty;$$

(4) y Codes

Avail and/or  
Special



Dist

A-1

(II)  $\sup_n \text{var}(X_n) < \infty$  and there exists  $\epsilon > 0$  such that

$$\sum_{i=1}^n \mu^{1/2}(i) = O(n/(\log n)^{1+\epsilon}); \quad (5)$$

(III) (4) holds,  $X_1, X_2, \dots$  are identically distributed and  $E|X_1| < \infty$  (the existence of variance is not assumed). Then (3) is true.

*Remarks:*

1. Part (I) of Theorem 2 can be compared with a result of Blum *et al* [1], who assumes that  $\{X_n\}$  is a  $\star$ -mixing sequence instead of (4). Note that this assumption does not follow from (4). We can easily construct a pairwise independent sequence which is not  $\star$ -mixing.

2. Parts (I) and (II) of Theorem 2 can also be compared with some results (see Theorem 3.7.2 and Theorem 3.7.4 of Stout [5]) derived from Serfling [4]. The conditions of these results involve correlation coefficients between two variables in the sequence.

3. Part (III) of Theorem 2 extends Theorem 1 of Etemadi [2]. The assumption that  $\{X_i\}$  is identically distributed can be somewhat relaxed, for example, it can be replaced by the condition that there exists a random variable  $Y$  such that  $P(|X_n| \geq x) \leq P(|Y| \geq x)$  for all  $n \geq 1$  and  $x \geq 0$ . We also mention a related result of Blum *et al* [1] Theorem 1. They assume that  $\{X_n\}$  is identically distributed, the distribution of  $X_1$  has a moment generating function in the neighborhood of zero and that  $\{X_n\}$  is  $\star$ -mixing. Under these more stronger conditions they prove that  $P(|S_n - ES_n|/n \geq \epsilon)$  tends to zero exponentially.

## 2. PROOF OF THE THEOREMS

In deducing our results we shall borrow a trick from Etemadi [2].

The following well-known facts concerning  $\alpha(X,Y)$  and  $\beta(X,Y)$  will be used:

$$|\text{cov}(X,Y)| \leq 10(\alpha(X,Y))^{\delta/(2+\delta)}(E|X|^{2+\delta}E|Y|^{2+\delta})^{1/(2+\delta)}, \quad \delta > 0 \quad (6)$$

$$|\text{cov}(X,Y)| \leq 2(\beta(X,Y)\text{var}(X)\text{var}(Y))^{1/2}. \quad (7)$$

For a proof, see Ibragimov and Linnik [3]. Also it is trivially true that

$$\alpha(XI_C(X), YI_D(Y)) \leq \alpha(X,Y), \quad \beta(XI_C(X), YI_D(Y)) \leq \beta(X,Y) \quad (8)$$

$$\alpha(X-a, Y-b) = \alpha(X,Y), \quad \beta(X-a, Y-b) = \beta(X,Y), \quad (9)$$

where  $C$  and  $D$  are Borel sets in  $R^1$  and  $a, b$  are constants.

*Proof of Theorem 1.* In view of (9), by defining  $X_n^+ = X_n I(X_n > 0)$ ,  $X_n^- = -X_n I(X_n \leq 0)$ ,  $n \geq 1$ , we can assume without loss of generality that  $X_n \geq 0$ ,  $n \geq 1$ . Define

$$Y_n = (X_n - EX_n)I(|X_n - EX_n| < n^{1/p+\epsilon_1}), \quad n \geq 1, \quad (10)$$

$$S_n^* = \sum_{i=1}^n (Y_i - EY_i),$$

where  $\epsilon_1 > 0$  is a constant to be chosen later.

From condition (1) we have  $\sum_{n=1}^{\infty} P(X_n - EX_n \neq Y_n) < \infty$  and  $\lim_{n \rightarrow \infty} EY_n = 0$ .

Therefore, (3) is equivalent to

$$\lim_{n \rightarrow \infty} S_n^*/n = 0, \quad \text{a.s.} \quad (11)$$

Now fix  $\alpha > 1$  and let  $k_n = [\alpha^n]$ . For positive integer  $m$  sufficiently large, there exists  $n$  such that  $k_n \leq m < k_{n+1}$ , and  $n \rightarrow \infty$  as  $m \rightarrow \infty$ . From (1) we have

$$\sup_n E|Y_n| \equiv C < \infty. \quad (12)$$

Here and in the sequel  $C$  is an unimportant constant which is allowed to change. Since  $Y_n \geq 0$ , it follows that

$$\begin{aligned} S_m^* - S_{k_n}^* &\geq -(m - k_n)C, \quad \text{when } S_m^* < S_{k_n}^*, \\ S_m^* - S_{k_n}^* &\leq S_{k_{n+1}}^* - S_{k_n}^* + (k_{n+1} - m)C, \quad \text{when } S_m^* \geq S_{k_n}^*. \end{aligned}$$

Hence

$$|S_m^*/m - S_{k_n}^*/k_n| \leq \left| \frac{k_{n+1}}{k_n} \frac{S_{k_{n+1}}^*}{k_{n+1}} - \frac{S_{k_n}^*}{k_n} \right| + \frac{k_{n+1} - k_n}{k_n} C. \quad (13)$$

From (13) it follows that if we have shown that

$$\lim_{n \rightarrow \infty} S_{k_n}^*/k_n = 0, \quad \text{a.s.} \quad (14)$$

Then we would have

$$\limsup_{m \rightarrow \infty} |S_m^*/m| \leq (\alpha - 1)C, \quad \text{a.s.}$$

For any  $\alpha > 1$ , hence (11).

By Borel-Cantelli lemma, in order to prove (14), we have only to show that

$$\sum_{n=1}^{\infty} \text{var}(S_{k_n}^*)/k_n^2 < \infty. \quad (15)$$

By (6), (8) and (9), we have for any  $\delta > 0$ :

$$\begin{aligned} \text{Var}(S_{k_n}^*) &= \sum_{i,j=1}^{k_n} \text{cov}(Y_i, Y_j) \\ &\leq C \sum_{i,j=1}^{k_n} (\alpha(X_i, X_j))^{\delta/(2+\delta)} (E|Y_i|^{2+\delta} E|Y_j|^{2+\delta})^{1/(2+\delta)}. \end{aligned} \quad (16)$$

From (1) it follows that

$$E|Y_n|^{2+\delta} \leq Cn^{(2+\delta-p)(1/p+\epsilon_1)}, \quad n = 1, 2, \dots \quad (17)$$

First consider the case  $p > 2$ . From (2), (16) and (17) we obtain

$$\begin{aligned} \text{var}(S_{k_n}^*) &\leq C \sum_{i,j=1}^{k_n} (\alpha(X_i, X_j))^{\delta/(2+\delta)} (ij)^{(2+\delta-p)(1/p+\epsilon_1)/(2+\delta)} \\ &\leq C \sum_{i,j=1}^{k_n} (\alpha(X_i, X_j))^{\delta/(2+\delta)} i^{2(2+\delta-p)(1/p+\epsilon_1)/(2+\delta)} \\ &\leq C \sum_{i,j=1}^{k_n} i^{-(2/p+\epsilon)\delta/(2+\delta)} \sum_{i=1}^{k_n} i^{2(2+\delta-p)(1/p+\epsilon_1)/(2+\delta)}. \end{aligned} \quad (18)$$

Noticing  $2/p < 1$ , we can assume that  $2/p + \epsilon < 1$ . Hence from (18) we have

$$\text{var}(S_{k_n}^*) \leq Ck_n^{-\frac{(2/p+\epsilon)\delta}{(2+\delta)} + \frac{2(2+\delta-p)(1/p+\epsilon_1)}{(2+\delta)} + 2}. \quad (19)$$

This inequality holds for any  $\delta > 0$ . Now we choose  $\epsilon_1 \in (0, \epsilon/2)$ , then

$$\lim_{\delta \rightarrow \infty} \{-\frac{(2/p+\epsilon)\delta}{(2+\delta)} + \frac{2(2+\delta-p)(1/p+\epsilon_1)}{(2+\delta)}\} = -\epsilon + 2\epsilon_1 \equiv \eta < 0.$$

Therefore, choosing  $\delta$  sufficiently large, from (19) we obtain

$$\text{var}(S_{k_n}^*) \leq Ck_n^{2-\eta}. \quad \text{Hence (15) is true in view of } \sum_{n=1}^{\infty} k_n^{-\eta} < \infty.$$

Next assume that  $p = 2$ . Again, choose  $\epsilon_1 \in (0, \epsilon/2)$ . Choose  $\delta > 0$  sufficiently small, such that  $(1+\epsilon)\delta/(2+\delta) < 1$ . We still have (19), with  $p = 2$ . Since

$$-(1+\epsilon)\delta/(2+\delta) + 2\delta(1/2+\epsilon_1)/(2+\delta) = -(\epsilon - 2\epsilon_1)\delta/(2+\delta) < 0,$$

(15) holds again.

Finally, consider the case  $1 < p < 2$ . In this case we have, instead of (18),

$$\text{var}(S_{k_n}^*) \leq C \sum_{i=1}^{k_n} i^{-\left(\frac{p}{2p-2} + \epsilon\right)\delta/(2+\delta)} \sum_{i=1}^{k_n} i^{2(2+\delta-p)(1/p + \epsilon_1)/(2+\delta)}. \quad (20)$$

Write  $\delta_0 = 2(p/(2p-2) - 1 + \epsilon)^{-1}$ . Since  $1 < p < 2$ , we have  $\delta_0 > 0$ . Choose  $\epsilon_1 > 0$  sufficiently small, such that

$$0 < \delta < \delta_0 \Rightarrow 2(2+\delta-p)(1/p + \epsilon_1)/(2+\delta) \leq 1 - \eta$$

where  $\eta > 0$  does not depend on  $\delta$ , as long as  $0 < \delta < \delta_0$ . Because  $(p/(2p-2) + \epsilon)\delta/(2+\delta) < 1$  for  $0 < \delta < \delta_0$  and  $(p/(2p-2) + \epsilon)\delta_0/(2+\delta_0) = 1$ , one can find  $\delta \in (0, \delta_0)$ , such that

$$1 - \eta/2 < (p/(2p-2) + \epsilon)\delta/(2+\delta) < 1.$$

For this  $\delta$  we have, by (20),

$$\text{var}(S_{k_n}^*) \leq C k_n^{-(1-\eta/2) + 1 + (1-\eta) + 1} \leq C k_n^{-\eta/2}.$$

So we obtain (15) again. Theorem 1 is proved.

*Proof of Theorem 2. Part (I):* Again we can assume  $X_n \geq 0$ . Write  $Y_n = X_n - EX_n$  and  $S_n^* = \sum_{i=1}^n Y_i$ . From  $\sup E|X_n| < \infty$  we have  $\sup E|Y_n| < \infty$ . Using the same argument employed in proving Theorem 1, we reduce the proof of (11) to that of (15). From (4), (7) and (9),

$$\begin{aligned} \sum_{n=1}^{\infty} \text{var}(S_{k_n}^*)/k_n^2 &= \sum_{n=1}^{\infty} k_n^{-2} \sum_{i,j=1}^{k_n} \text{cov}(Y_i, Y_j) \\ &\leq C \sum_{n=1}^{\infty} k_n^{-2} \sum_{i,j=1}^{k_n} (\mu(|i-j|) \text{var}(X_i) \text{var}(X_j))^{1/2} \\ &\leq C \sum_{n=1}^{\infty} k_n^{-2} \sum_{i=0}^{k_n} \mu^{1/2}(i) \sum_{i=1}^{k_n} \text{var}(X_i) \\ &\leq C \sum_{n=1}^{\infty} k_n^{-2} \sum_{i=1}^{k_n} \text{var}(X_i) \end{aligned} \quad (21)$$

$$\begin{aligned} &\leq C \sum_{n=1}^{\infty} \text{var}(X_n)/n^2 < \infty. \end{aligned} \quad (22)$$

Part (II) is proved in much the same way as Part (I), only that we replace  $Ck_n$  for  $\sum_{i=1}^{k_n} \text{var}(X_i)$  and  $Ck_n/(\log n)^{1+\epsilon}$  for  $\sum_{i=1}^{k_n} \mu^{1/2}(i)$  in (21) to obtain (22). Part (III) is proved by truncating  $X_n$  at  $n$  and combining the reasoning above and that of Etemadi [2].

### 3. AN EXAMPLE

Consider the autoregression model

$$X_n = a_1 X_{n-1} + \dots + a_m X_{n-m} + e_n, \quad n = 0, \pm 1, \pm 2, \dots \quad (23)$$

We want to show that under certain conditions it is true that

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n X_i / n = 0, \quad \text{a.s.} \quad (24)$$

for any solution of (23). Suppose that the following conditions are satisfied:

1.  $\{e_n, n = 0, \pm 1, \dots\}$  is a sequence of independent real random variables, and

$$Ee_n = 0, \quad n = 0, \pm 1, \dots, \quad \sup_{-\infty < n < \infty} E|e_n|^p = C < \infty \quad \text{for some } p > 1. \quad (25)$$

where, as before,  $C$  is an unimportant constant which is allowed to change.

2.  $e_n$  has a density  $f_n$  satisfying the Lipschitz condition over  $R'$ :

$$|f_n(x) - f_n(y)| \leq C|x - y|, \quad n = 0, \pm 1, \pm 2, \dots \quad (26)$$

where  $C$  does not depend on  $n$ .

3.  $a_1, a_2, \dots, a_m$  are real constants, and the equation  $1 - a_1 z - \dots - a_m z^m = 0$  has all its root outside the unit circle.

Under the condition 1 and 3, the general real solution of (23) has

the form

$$X_n = \sum_{t=0}^{\infty} b_t e_{n-t} + \sum_{j=0}^J \rho_j^n \sum_{\ell=0}^{m_j-1} n^{\ell} (\xi_{j\ell} \cos n\omega_j + \eta_{j\ell} \sin n\omega_j) \equiv \tilde{X}_n + X_n^* \quad (27)$$

where  $b_0 = 1$ ,  $b_2, b_3, \dots$  are real constants such that

$$|b_t| \leq CH^t, \quad t = 0, 1, 2, \dots \quad \text{for some } H \in (0, 1). \quad (28)$$

$\rho_j$  and  $\omega_j$ ,  $j = 1, \dots, J$ , are real constants,  $0 < \rho_j < 1$ ,  $j = 1, \dots, J$ ,  $m_1 + \dots + m_J = m$ , and  $\xi_{j\ell}, \eta_{j\ell}$ ,  $\ell = 1, \dots, m_j$ ,  $j = 1, \dots, J$ , are arbitrary random variables. From (25), (27) and (28) it follows that

$$E\tilde{X}_n = 0, \quad n = 0, 1, 2, \dots, \quad \sup_{-\infty < n < \infty} E|\tilde{X}_n|^p = C < \infty. \quad (29)$$

Let  $n, N$  be positive integers,  $n < N$ . Define

$$Y_{nN} = \sum_{t=0}^{N-n-1} b_t e_{N-t}, \quad Z_{nN} = \sum_{t=N-n}^{\infty} b_t e_{N-t}.$$

Since  $b_0 = 1$ , from (26) it follows that the density  $g_{nN}$  of  $Y_{nN}$  obeys Lipschitz's condition with the same constant  $C$  as in (26). Also

$$\sup\{E|Y_{nN}|^p : 1 \leq n < N < \infty\} = C < \infty. \quad (30)$$

Now let  $q_1$  be a positive constant,  $q_2 = 2q_1$ . Define the event

$$D_{nN} = \{|Z_{nN}| \geq (N-n)^{-q_2}\}. \quad (31)$$

(25) entails  $\sup_{-\infty < n < \infty} E|e_n| = C < \infty$ . Hence

$$P(D_{nN}) \leq C(N-n)^{q_2} \sum_{t=N-n}^{\infty} H^t \leq C(N-n)^{q_2} H^{N-n}. \quad (32)$$

Let  $G$  be a Borel set in  $R'$ ,  $h$  be a constant.  $G - h$  is defined as the

set  $\{g-h: g \in H\}$ . Write  $\tilde{G} = G \cap \{u: |u| \leq (N-n)^{q_1}\}$ ,  $G^* = G \setminus \tilde{G}$ . If  $|h| < 1$ , we have

$$\begin{aligned}
 & |P(Y_{nN} \in G) - P(Y_{nN} \in G-h)| \\
 & \leq |P(Y_{nN} \in \tilde{G}) - P(Y_{nN} \in \tilde{G}-h)| + P(Y_{nN} \in G^*) + P(Y_{nN} \in G^*-h) \\
 & \leq \int_{\tilde{G}} |g_{nN}(u) - g_{nN}(u-h)| du + P(|Y_{nN}| > (N-n)^{q_1}) + P(|Y_{nN}| > (N-n)^{q_1} - 1) \\
 & \leq C(N-n)^{q_1} h + C(N-n)^{-q_1} + C[(N-n)^{q_1} - 1]^{-1} \\
 & \leq C(N-n)^{q_1} h + C(N-n)^{-q_1}.
 \end{aligned} \tag{33}$$

Now let  $A$  and  $B$  be two Borel sets in  $R'$ . We proceed to estimate

$|P(\tilde{X}_n \in A, \tilde{X}_N \in B) - P(\tilde{X}_n \in A)P(\tilde{X}_N \in B)|$ . From (32), (33) and the independence of  $e_1, e_2, \dots$ , we have

$$\begin{aligned}
 |P(\tilde{X}_n \in B | e_n, e_{n-1}, \dots) - P(Y_{nN} \in B)| &= |P(Y_{nN} \in B - Z_{nN} | Z_{nN}) - P(Y_{nN} \in B)| \\
 &\leq C(N-n)^{-(q_2-q_1)} + C(N-n)^{-q_1} \\
 &\leq C(N-n)^{-q_1},
 \end{aligned} \tag{34}$$

when  $D_{nN}$  does not occur. But

$$\begin{aligned}
 |P(\tilde{X}_N \in B) - P(Y_{nN} \in B)| &= |P(Y_{nN} \in B - Z_{nN}) - P(Y_{nN} \in B)| \\
 &= |P(D_{nN}^C)P(Y_{nN} \in B - Z_{nN}) + P(D_{nN})P(Y_{nN} \in B - Z_{nN} | D_{nN}) \\
 &\quad - P(Y_{nN} \in B)| \\
 &\leq P(D_{nN}) + |P(Y_{nN} \in B - Z_{nN} | D_{nN}^C) - P(Y_{nN} \in B)| + P(D_{nN}) \\
 &\leq 2P(D_{nN}) + C(N-n)^{-q_1} \leq C(N-n)^{q_2} H^{N-n} + C(N-n)^{-q_1} \\
 &\leq C(N-n)^{-q_1}.
 \end{aligned} \tag{35}$$

From (34) and (35) we get

$$|P(\tilde{X}_N \in B | e_n, e_{n-1}, \dots) - P(\tilde{X}_N \in B)| \leq C(N-n)^{-q_1}$$

when  $D_{nN}$  does not occur. If  $P(\tilde{X}_n \in B) \geq C(N-n)^{-q_1}$ , then from (33) and (35) we obtain

$$P(\tilde{X}_n \in A, \tilde{X}_N \in B) \geq [P(\tilde{X}_N \in B) - C(N-n)^{-q_1}][P(\tilde{X}_n \in A) - C(N-n)^{q_2} H^{N-n}]. \quad (36)$$

Also

$$P(\tilde{X}_n \in A, X_N \in B) \leq [P(\tilde{X}_N \in B) + C(N-n)^{-q_1}][P(\tilde{X}_n \in A) + C(N-n)^{q_2} H^{N-n}]. \quad (37)$$

From (36) and (37) we have

$$\begin{aligned} |P(\tilde{X}_n \in A, \tilde{X}_N \in B) - P(\tilde{X}_n \in A)P(\tilde{X}_N \in B)| &\leq C(N-n)^{-q_1} + C(N-n)^{q_2} H^{N-n} \\ &\quad + C(N-n)^{q_1} H^{N-n} \leq C(N-n)^{-q_1}, \end{aligned} \quad (38)$$

where  $C$  does not depend on  $A, B$ . (38) is proved when  $P(\tilde{X}_n \in B) \geq C(N-n)^{-q_1}$ . If  $P(\tilde{X}_N \in B) < C(N-n)^{-q_1}$ , (38) is trivially true. Therefore we get

$$\alpha(\tilde{X}_n, \tilde{X}_N) \leq C(N-n)^{-q_1}. \quad (39)$$

Now choose  $q_1 = p/(2p-2) + 2$ . From (39) we see that the condition (2) is satisfied. This, together with (29), gives, by Theorem 1,

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n \tilde{X}_i / n = 0, \quad \text{a.s.} \quad (40)$$

From the expression of  $X_n^*$ , it is readily seen that

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n X_i^* / n = 0, \quad \text{a.s.} \quad (41)$$

From (27), (40) and (41), we obtain (24).

The conclusion (40) does not follow from the ergodic theorem of stationary process, since  $\{e_n\}$  is not assumed to be identically distributed, so  $\{X_n\}$  may not be a strictly stationary process.

#### REFERENCES

- [1] BLUM, J.R., HANSON, D.L. and KOOPMANS, L.H. (1963). On the strong law of large numbers for a class of stochastic process. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* 2, 1-11.
- [2] ETEMADI, N. (1981). An elementary proof of strong law of large numbers. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* 55, 119-122.
- [3] IBRAGIMOV, I.A. and LINNIK, J.V. (1965). *Independent and Stationary Connected Variables* (Nauka, Moscow), English translation (Noordhoff, Groningen, 1971).
- [4] SERFLING, R.J. (1970). Convergence properties of  $S_n$  under moment restrictions. *Ann. Math. Statist.* 41, 1235-1248.
- [5] STOUT, W.F. (1974). *Almost Sure Convergence*. Academic Press.

END

DATE

FILMED

7-88

Dtic